

# Medial Patellofemoral Ligament Repair and Reconstruction

The knee consists of four bones that form three joints. The femur is the large bone in the thigh and attaches by ligaments and a capsule to the tibia, the large bone in the lower leg commonly referred to as the shin bone. Next to the tibia is the fibula, which runs parallel to the tibia on the outside of the leg. The patella, commonly called the kneecap, is embedded in the quadriceps and patellar tendon which articulates with the front of the femur, which forms the patellofemoral joint. The patella acts as a pulley to increase the amount of force that the quadriceps muscle can generate.<sup>1</sup> The patella sits in a groove on the end of the femur called the trochlear groove. This groove varies in depth from person to person. When the knee bends, the patella travels down the groove and as the knee straightens the patella moves up the groove. As the patella travels up and down in the trochlear groove, the patella should maintain congruent bony alignment, which is often referred to as normal patellar tracking.

There are several structures that work together to keep the patella aligned and stabilized in the femoral groove to prevent the patella from excessive lateral movement (movement towards the outside of the leg). The lateral aspect of the trochlear groove is normally about 1 centimeter higher than the medial (inside of the leg) aspect of the trochlear groove, which helps keep the patella in the trochlear groove by providing a buttress on the lateral side (Figure 1).<sup>2</sup> This provides the main resistance to lateral patellar translation (which is the most common direction of displacement), especially beyond 20 degrees of knee flexion.<sup>3</sup>

People who have a shallow trochlea are more susceptible to patellar instability.

Proper stabilization of the patella is also affected by the soft tissue structures (ligaments and muscles) surrounding the knee. The medial patellofemoral ligament (MPFL) is a continuation of the deep retinaculum and vastus medialis oblique (VMO) muscle fibers (inner portion of the quadriceps muscle) on the inside of the knee. These structures provide a significant force (near 60% total) against lateral displacement of the patella, as their force is directed inward or medially.<sup>2,4</sup> The MPFL is the primary restraint to lateral displacement of the patella during the first 20 to 30 degrees of knee flexion.<sup>3</sup> This ligament is a passive stabilizer and extends from the upper inner side of the patella to the medial aspect of the femur. The patellomeniscal ligament and retinaculum also contribute more than 20% of the restraining force.



**Figure 1.** Radiograph of the patellofemoral joint with the knee in slight flexion. The lateral aspect of the trochlear groove is normally about 1 cm higher than the medial.

These ligaments can be injured and torn with an initial acute traumatic patellar dislocation (kneecap quickly going out of place). The most common mechanism for a patellar dislocation is a forceful inward rotation of the body on a planted foot. This radiograph is of a 12-year-old boy in the emergency room after such an injury (Figure 2) Often the patella will go back in to place (or relocate to the groove) as the knee is gently straightened. In this case the patient was unable to straighten his knee in the emergency room and his patella was still dislocated laterally. Note on the radiograph that there is no overlap of the femur and patella.

In studying 26 patients who had an acute patellar dislocation at a mean age of 18 years, Nomura et al. reported evidence of MPFL damage in 96% (26/27) of the knees examined during open surgical exploration). In an acute patellar dislocation, when a tear of the MPFL is identified, surgical repair (fixing the original ligament) of the MPFL may be a good treatment option. In the young athletic population, recurrence rates for patients treated conservatively are high with some studies reporting 40%. In recurrent or chronic patellar dislocations, it may be necessary to perform reconstruction of the MPFL. Reconstruction differs from repair in that graft tissue (such as a hamstring tendon) is used to replace or reinforce the MPFL. The MPFL reconstruction may also be combined with other patellar stabilization procedures.

A quality post-operative rehabilitation program is essential to having a successful outcome from a MPFL procedure. The goals of rehabilitation will initially focus on protection for healing, mobility and range of motion.

After this early phase, strengthening and neuromuscular control is emphasized throughout the entire leg and core. In the final stages of rehabilitation, the focus will be on dynamic lower extremity control during sport specific movements, such as change of direction and rotational movements.

The UW Health Sports Medicine rehabilitation guidelines below are presented in a criterion-based progression. General time frames are given for reference to the average, but individual patients will progress at different rates depending on their age, associated injuries, pre-injury health status, rehabilitation compliance and injury severity. Modifications in the specific time frames, restrictions and precautions may also be made to protect healing tissues based on the specific surgical repair/ reconstruction procedure performed.




Figure 2. Radiograph of the knee, arrows show the laterally dislocated patella

Phase I (surgery to 6 weeks after surgery)	
<b>Rehabilitation appointments</b>	<ul style="list-style-type: none"> <li>• Rehabilitation appointments to begin 2-3 days after surgery and to continue 1-2 times per week</li> </ul>
<b>Rehabilitation goals</b>	<ul style="list-style-type: none"> <li>• Protect surgical knee</li> <li>• Restore normal knee range of motion</li> <li>• Full passive knee extension should be achieved within first two weeks, though these patients do not often struggle with extension</li> <li>• Goal of a 90 degrees knee flexion should be achieved by 6 weeks</li> <li>• Normalize gait</li> <li>• Eliminate effusion (swelling)</li> <li>• Restore leg control and quadriceps activation</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>• Weight bearing as tolerated on crutches with post-operative brace locked in extension.</li> <li>• Begin ambulation with 2 crutches, then progress to 1 crutch and then no crutches once patient shows sufficient quad control and gait mechanics are normalized</li> <li>• Avoid rotational movements through knee and lower extremity</li> </ul>
<b>Suggested therapeutic exercise</b>	<ul style="list-style-type: none"> <li>• Gait drills with emphasis on symmetrical loading, heel strike at initial contact, appropriate quad activation during stance (avoid hyperflexed or hyperextended knee during mid-stance), and adequate push-off</li> <li>• Range of motion (ROM): (pearl: provide manual lateral patellar stabilization when first initiating range of motion)             <ul style="list-style-type: none"> <li>• Knee extension with foot propped on bolster</li> <li>• Heel slides</li> <li>• Knee flexion wall slides</li> <li>• Passive knee flexion over edge of plinth</li> <li>• Medial Patellar mobilizations</li> <li>• Prone knee flexion</li> </ul> </li> <li>• Strengthening:             <p><b>Note: it is recommended that all quadriceps strengthening be performed in conjunction with neuromuscular electrical stimulation (NMES). Please see Appendix for recommendations on NMES units for patients, parameters for use, and treatment recommendations for NMES during quadriceps strengthening.</b></p> <ul style="list-style-type: none"> <li>• Quadriceps, hamstring, and gluteal sets</li> </ul> </li> </ul>

	<ul style="list-style-type: none"> <li>• Four-way leg lifts in standing or lying down positions for hip strengthening</li> <li>• Bridging</li> <li>• Ankle isotonic with resistance band</li> <li>• Weight shifting drills</li> <li>• Heel raises</li> <li>• Balance drills beginning with double leg and progressing to single leg</li> <li>• Trunk stability work             <ul style="list-style-type: none"> <li>• Supine core and transverse abdominus activation with upper and/or lower extremity movement</li> <li>• Anti-rotation press variations in stable lower extremity positions</li> <li>• Suitcase carry with gait and/or marching drills</li> </ul> </li> </ul>
<b>Cardiovascular exercise</b>	<ul style="list-style-type: none"> <li>• Upper body circuit training or use of an upper body ergometer</li> </ul>
<b>Progression criteria</b>	<ul style="list-style-type: none"> <li>• Non-painful knee flexion AROM to 90 degrees</li> <li>• Full weight bearing with normalized gait mechanics without the use of assistive device</li> <li>• Single leg balance for 15 seconds with good control</li> </ul>

<b>Phase II (begin after meeting Phase I criteria, usually 6 weeks after surgery)</b>	
<b>Rehabilitation appointments</b>	<ul style="list-style-type: none"> <li>Rehabilitation appointments are 1-2 times per week</li> </ul>
<b>Rehabilitation goals</b>	<ul style="list-style-type: none"> <li>Maintain/fully restore normal knee range of motion</li> <li>Maintain protection of post-surgical knee</li> <li>Increase functional activity</li> <li>Good limb control and no pain with functional movements</li> <li>First Progressive Testing session should occur at 12 weeks after surgery:</li> <li>LSI &gt;90% on single leg press and Y-Balance</li> <li>Quadriceps strength deficit of &lt;30% on Biodex strength test</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>Use of lateral buttress knee sleeve if directed by physician or physical therapist</li> <li>Avoid over-stressing fixation of graft – continued caution with rotational movement                             <ul style="list-style-type: none"> <li>For patients with patellar or trochlear chondroplasty or cartilage defect, avoid excessive or abnormal patellofemoral joint stresses during open and closed chain strengthening</li> </ul> </li> <li>Avoid post-activity swelling</li> <li>Full ROM into flexion and extension should be achieved if patient has not already regained full ROM</li> <li>No impact</li> </ul>
<b>Suggested therapeutic exercise</b>	<ul style="list-style-type: none"> <li>Range of motion work as needed</li> <li>Continue NMES</li> <li>Continued functional closed kinetic chain strengthening with caution to avoid dynamic valgus or medial knee displacement:</li> <li>Progressions of double leg squats</li> <li>Split stance work progressing into lunge variations</li> <li>Single leg bridging</li> <li>Controlled single leg squats</li> <li>Progression of weight on leg press and progression to single leg press</li> <li>Multidirectional band walks</li> <li>Continued hamstring and gluteal strengthening</li> <li>Continued trunk strength and stability work</li> </ul>
<b>Cardiovascular exercise</b>	<ul style="list-style-type: none"> <li>Upper body circuit training or use of an upper body ergometer</li> <li>Stationary bike</li> </ul>

<b>Progression criteria</b>	<ul style="list-style-type: none"><li>• Normal gait on level surfaces</li><li>• LSI &gt;90% on single leg press and Y-Balance</li><li>• Quadriceps strength deficit of &lt;30% on Biodex strength test</li><li>• At least 12 weeks after surgery</li></ul>
-----------------------------	--

<b>Phase III begin after meeting Phase II criteria, usually 12-16 weeks after surgery)</b>	
<b>Rehabilitation appointments</b>	<ul style="list-style-type: none"> <li>• Rehabilitation appointments once every 1-2 weeks</li> </ul>
<b>Rehabilitation goals</b>	<ul style="list-style-type: none"> <li>• No effusion (swelling)</li> <li>• Return to full functional activities</li> <li>• Improve quadriceps strength</li> <li>• Improve hip and trunk strength</li> <li>• Improve balance and proprioception</li> <li>• Improve patient confidence and mental readiness to return to higher level movement patterns and eventually return to sport</li> </ul>
<b>Suggested therapeutic exercises</b>	<ul style="list-style-type: none"> <li>• Continue closed chain strengthening beginning with single plane and progressing to multi-planar</li> <li>• Progression of hip and core strengthening</li> <li>• Progression of speed during strengthening drills to increase rate of force development (RFD) and for impact preparation</li> <li>• Initiate low amplitude agility drills in the sagittal and frontal plane – caution with frontal plane and avoid transverse plane initially because of potential for dynamic valgus</li> <li>• Impact control exercises in sagittal and frontal plane beginning 2 feet to 2 feet, progressing toward 1 foot to the other foot (bounding)                             <ul style="list-style-type: none"> <li>• Initiate return to running progression once patient shows good single leg control and tolerance to bounding</li> </ul> </li> <li>• Stretching for patient specific muscle imbalances</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>• Avoid post-activity swelling</li> </ul>
<b>Cardiovascular exercise</b>	<ul style="list-style-type: none"> <li>• Stationary bike</li> <li>• Treadmill walking</li> <li>• Swimming with flutter kick (avoid breaststroke kick)</li> <li>• Stair Master</li> <li>• Elliptical</li> </ul>
<b>Progression criteria</b>	<ul style="list-style-type: none"> <li>• No effusion (swelling)</li> <li>• No patellar apprehension</li> <li>• Good control and no pain with squats and lunges and impact drills</li> <li>• Quadriceps strength deficit of &lt;15% on Biodex strength test</li> </ul>


<b>Phase IV (begin after meeting Phase III criteria, usually 20 weeks after surgery)</b>	
<b>Rehabilitation appointments</b>	<ul style="list-style-type: none"> <li>Rehabilitation appointments once every 1-2 weeks</li> </ul>
<b>Rehabilitation goals</b>	<ul style="list-style-type: none"> <li>Good eccentric and concentric multi-planar dynamic neuromuscular control (including impact) to allow for return to work/sports</li> </ul>
<b>Suggested therapeutic exercises</b>	<ul style="list-style-type: none"> <li>Progression of impact control exercises to 1 foot to the same foot (hopping)</li> <li>Movement control exercises beginning with low velocity, single plane activities and progressing to higher velocity, multi-planar activities</li> <li>Progression to multi-planar agility drills with progressively increasing velocity and amplitude</li> <li>Sport/work specific balance and proprioception drills</li> <li>Continue lower extremity and trunk strengthening</li> <li>Stretching for patient specific muscle imbalances</li> </ul>
<b>Precautions</b>	<ul style="list-style-type: none"> <li>Loaded horizontal abduction (bench press, push ups, etc) should begin and progress in a very gradual manner. Bench should include protecting the range and limiting to a maximum of 50% of pre-injury max for the first 18-22 weeks.</li> </ul>
<b>Cardiovascular exercise</b>	<ul style="list-style-type: none"> <li>Replicate sport or work specific energy demands</li> </ul>
<b>Progression criteria</b>	<ul style="list-style-type: none"> <li>Return to sport/work criteria:</li> <li>Quadriceps strength deficit of &lt;10% on Biodex strength test</li> <li>LSI of &gt;90% on jump testing and all four functional hop tests</li> <li>Dynamic neuromuscular control with multi-planar activities and without pain, instability, or swelling</li> <li>Patient confidence to return to sport</li> <li>Approval from the physician and sports rehabilitation provider</li> </ul>



Appendix

NMES Treatment Parameters:

- Amplitude/Intensity (mA): as high as patient can tolerate
- Pulse Width ( $\mu$ s): 100-400  $\mu$ s
- Pulse Rate (Hz): 50-100 Hz
- Frequency: NMES should be dosed the same way you would dose quadriceps strengthening exercise; early in the rehabilitation process, this should be used every day or even multiple times a day. As the patient progresses and the intensity of their exercise increases, the frequency will likely change to 2-4 times a week. While the exact treatment plan will vary with every patient, the diagram below (Figure 3) from Spector et al. outlines a proposed algorithm for the frequency of quadriceps strengthening with NMES following knee surgery. The algorithm is further outlined below:
  - Quadriceps NMES to begin as early as possible after surgery with a high-volume approach (1 or more times per day).
  - After 1 week of treatment, assess patient’s response to NMES to gauge effectiveness – Fitzgerald criteria is recommended as a clinically feasible way to assess the patient’s response
  - If patient is responding well, continue treatment for 2 more weeks
  - Re-assess response to NMES – look for signs of activation failure:
    - Inability to perform straight leg raise without extensor lag
    - In ability to consistently perform quadriceps set with superior patellar glide
    - Patient reported difficulty with muscle control
  - If activation failure is present, continue treatment with high-volume
  - If activation failure is NOT present, progress to quadriceps NMES with a low-volume approach (approximately 3-6 times per week)



*Figure 3.* Recommended treatment algorithm from Spector et al. for application of NMES with quadriceps strengthening following knee surgery.

\*Fitzgerald criteria: NMES should produce full tetanic contraction of quadriceps with a visual and/or palpable superior patellar glide

References

7. Fisher B, Nyland J, Brand E, Curtin B. Medial patellofemoral ligament reconstruction for recurrent patellar dislocation: a systematic review including rehabilitation and return-to-sports efficacy. *Arthroscopy*. 2010 Oct;26(10):1384-94.
8. Neuman DA. *Kinesiology of the Musculoskeletal System: Foundations for Physical Rehabilitation*. 1st ed. St. Louis, MO: Mosby; 2002.
9. Minkowitz R, Inzerillo C, Sherman OH. Patella Instability. *Bull NYU Hosp Jt Dis*. 2007;65(4):280-93.
10. Nomura E, Horiuchi Y, Inoue M. Correlation of MR imaging findings and open exploration of medial patellofemoral ligament injuries in acute patellar dislocations. *Knee*. 2002 May;9(2):139-43.
11. Labanca L, Rocchi JE, Laudani L, et al. Neuromuscular Electrical Stimulation Superimposed on Movement Early after ACL Surgery. *Med Sci Sports Exerc*. 2018;50(3):407-416. doi:10.1249/MSS.0000000000001462
12. Krych AJ, O'Malley MP, Johnson NR, et al. Functional testing and return to sport following stabilization surgery for recurrent lateral patellar instability in competitive athletes. *Knee Surgery, Sport Traumatol Arthrosc*. 2018;26(3):711-718. doi:10.1007/s00167-016-4409-2
1. Zhang F, Wang J, Wang F. Comparison of the clinical effects of open and closed chain exercises after medial patellofemoral ligament reconstruction. *J Phys Ther Sci*. 2014;26(10):1557-1560. doi:10.1589/jpts.26.1557
2. Glaviano NR, Saliba S. Can the Use of Neuromuscular Electrical Stimulation Be Improved to Optimize Quadriceps Strengthening? *Sports Health*. 2016;8(1):79-85. doi:10.1177/1941738115618174
3. Spector P, Laufer Y, Elboim Gabyzon M, Kittelson A, Stevens Lapsley J, Maffiuletti NA. Neuromuscular electrical stimulation therapy to restore quadriceps muscle function in patients after orthopaedic surgery. *J Bone Jt Surg - Am Vol*. 2016;98(23):2017-2024. doi:10.2106/JBJS.16.00192
4. Zaman S, White A, Shi WJ, Freedman KB, Dodson CC. Return-to-Play Guidelines After Medial Patellofemoral Ligament Surgery for Recurrent Patellar Instability: A Systematic Review. *Am J Sports Med*. 2018;46(10):2530-2539. doi:10.1177/0363546517713663
5. Saper MG, Fantozzi P, Bompadre V, Racicot M, Schmale GA. Return-to-Sport Testing After Medial Patellofemoral Ligament Reconstruction in Adolescent Athletes. *Orthop J Sport Med*. 2019;7(3):1-7. doi:10.1177/2325967119828953
6. Lightsey HM, Wright ML, Trofa DP, Popkin CA, Ahmad CS, Redler LH. Rehabilitation variability following medial patellofemoral ligament reconstruction. *Phys Sportsmed*. 2018;46(4):441-448. doi:10.1080/00913847.2018.1487240

At UW Health, patients may have may receive direction or educational materials that vary from this information. This information is not intended to replace the care or advice given by your physician or health care provider. It is neither intended nor implied to be a substitute for professional advice. Call your health provider immediately if you think you may have a medical emergency. Always seek the advice of your physician or other qualified health provider prior to starting any new treatment or with any question you may have regarding a medical condition.